

CALT-68-2111, hep-ph/9704370

NONPERTURBATIVE EFFECTS IN INCLUSIVE $\bar{B} \rightarrow X_s \gamma$

ZOLTAN LIGETI

California Institute of Technology, Pasadena, CA 91125

Uncertainties in the theoretical prediction for the inclusive $\bar{B} \rightarrow X_s \gamma$ decay rate are examined. Certain nonperturbative effects involving a virtual $c\bar{c}$ loop, which are calculable using the operator product expansion, are discussed.

The inclusive $\bar{B} \rightarrow X_s \gamma$ decay is sensitive to physics beyond the standard model, and the photon spectrum can help us better understand nonperturbative effects in other B decays. As the CLEO data¹ excludes large deviations from the standard model, it is important to know the theoretical predictions as precisely as possible. With the completion of the next-to-leading order calculation², the theoretical uncertainty in perturbation theory is only about 10%.

The effective weak interaction Hamiltonian at a scale $\mu (\sim m_b)$ is $H_{\text{eff}} = -(4G_F/\sqrt{2}) V_{ts}^* V_{tb} \sum_{i=1}^8 C_i(\mu) O_i(\mu)$. Here $O_2 = (\bar{s}_{L\alpha} \gamma_\mu b_{L\beta}) (\bar{c}_{L\beta} \gamma^\mu c_{L\alpha})$, O_1 differs from O_2 in the contraction of color indices, $O_3 - O_6$ are other four-quark operators, and $O_7 = (e/16\pi^2) m_b \bar{s}_L \sigma^{\mu\nu} F_{\mu\nu} b_R$.

For large enough photon energies, the matrix element of $C_7 O_7$ dominates the $\bar{B} \rightarrow X_s \gamma$ rate. Since $m_b \gg \Lambda_{\text{QCD}}$, this contribution is calculable by performing an operator product expansion (OPE) for the time ordered product³

$$T_{77} = \frac{i}{2m_B} \int d^4x e^{-iq \cdot x} \langle \bar{B}(v) | T \{ O_7^{\mu\dagger}(x) O_7^\nu(0) \} | \bar{B}(v) \rangle g_{\mu\nu}. \quad (1)$$

Here $O_7^\mu = (i e/8\pi^2) m_b \bar{s}_L \sigma^{\mu\lambda} q_\lambda b_R$. At fixed $q^2 = 0$, T_{77} has cuts in the complex $v \cdot q$ plane along $v \cdot q < m_b/2$ and $v \cdot q > 3m_b/2$ corresponding to final hadronic states X_s and $X_{b\bar{b}s}$, respectively. The $\bar{B} \rightarrow X_s \gamma$ decay rate is given by the discontinuity across the cut in the region $0 < v \cdot q < m_b/2$,

$$\frac{d\Gamma}{dE_\gamma} = \frac{4G_F^2 |V_{ts}^* V_{tb}|^2 C_7^2}{\pi^2} E_\gamma \text{Im } T_{77}. \quad (2)$$

Since the cuts are well-separated, this contribution can be computed assuming local duality at the scale m_b ($m_{X_s} = m_B$ at $v \cdot q = 0$).

At leading order in the OPE, the dimension-three operator $\bar{b} \gamma_\mu b$ occurs. Its matrix element gives a calculable contribution proportional to $\delta(E_\gamma - m_b/2)$,

^aTalk given at the FCNC'97 workshop, February 19-21, 1997, Santa Monica.

which is equal to the free quark decay result. Higher dimension operators give terms proportional to derivatives of this delta function. The leading nonperturbative corrections suppressed by $\Lambda_{\text{QCD}}^2/m_b^2$ are quite small, about -3% ³. To justify retaining only the lowest dimension operators whose matrix elements are known, the photon energy must be averaged over a region $\Delta E_\gamma \gg \Lambda_{\text{QCD}}$, and cannot be restricted to be too close to its maximal (i.e., end-point) value. Currently this is a source of significant uncertainty, since the photon spectrum is only measured over a region about 500 MeV from the end-point¹.

When operators in H_{eff} other than O_7 are included, there are contributions from diagrams in which the photon couples to light quarks. Typically, the leading logarithms are calculable for such processes⁴, but there are uncalculable contributions suppressed by a logarithm (or equivalently by α_s , but not by a power of the scale of the process), which can only be estimated using the fragmentation functions $D_{q \rightarrow \gamma X}$ and $D_{g \rightarrow \gamma X}$ deduced from other experiments or from models. Perturbative computations indicate that the contribution of light quark loops⁵ and the effects related to decay functions of light partons into a photon⁶ are both very small for decays into hard photons.⁶ Assuming that these calculations provide correct order of magnitude estimates⁷, such not power suppressed effects constitute less than five percent uncertainty in the theoretical prediction for the $\bar{B} \rightarrow X_s \gamma$ decay rate.

Nonperturbative effects involving the photon coupling to the charm quark contain matrix elements of local operators⁸ only suppressed by $\Lambda_{\text{QCD}}^2/m_c^2$ rather than $\Lambda_{\text{QCD}}^2/m_b^2$. Such effects could be sizable, since $m_b^2/m_c^2 \sim 10$ and the $\Lambda_{\text{QCD}}^2/m_b^2$ corrections are 3%. For a sufficiently heavy charm quark, nonperturbative corrections to the interference of O_2 and O_7 can be computed from the discontinuity of the diagram in Fig. 1. Analogous diagrams with more gluons are suppressed by additional powers of Λ_{QCD}/m_c . Denote the photon and gluon momenta by q and k respectively. Working to all orders in $k \cdot q/m_c^2$ since $|q| \sim m_b$, but neglecting $k \cdot q/m_b^2$, $k^2/m_{c,b}^2$, and m_s/m_b , gives⁹

$$T_{27} = -\frac{1}{2m_B} \langle \bar{B}(v) | \bar{b} m_b \sigma^{\nu\rho} q_\rho \frac{m_b \not{v} - \not{q}}{(m_b v - q)^2 + i\epsilon} \gamma^\mu (1 - \gamma_5) I_{\mu\nu} b | \bar{B}(v) \rangle. \quad (3)$$

Here $I_{\mu\nu}$ is a complicated operator involving all powers of $q \cdot iD/m_c^2$,

$$I_{\mu\nu} = \left(\frac{e}{16\pi^2} \right)^2 \frac{2}{9m_c^2} \left[\sum_{n=0}^{\infty} \frac{3[(n+1)!]^2 2^{n+3}}{(2n+4)!} \left(\frac{-q \cdot iD}{m_c^2} \right)^n \right] \varepsilon_{\mu\nu\lambda\beta} q^\beta q_\eta g_s G^{\lambda\eta}. \quad (4)$$

⁹For soft photons these are important. Interference effects where the photon couples to a light quark and either to the charm quark or through O_7 are also small for hard photons.




Figure 1: Feynman diagram that gives rise to T_{27} in Eq. (3). Interchange of the photon and gluon couplings to the charm loop is understood.

The covariant derivatives, D , act on the gluon field $G^{\lambda\eta}$, so the matrix elements of these operators are determined by the spacetime dependence of the chromomagnetic field in the B meson. The contribution of T_{27} to the $\bar{B} \rightarrow X_s \gamma$ decay rate is given by Eq. (2) with $C_7^2 T_{77}$ replaced by $2C_2 C_7 T_{27}$.

For the leading $n = 0$ term in Eq. (4), the matrix element in Eq. (3) is known from the $B^* - B$ mass splitting. This gives

$$\frac{\delta\Gamma(\bar{B} \rightarrow X_s \gamma)}{\Gamma(\bar{B} \rightarrow X_s \gamma)} = -\frac{C_2}{9C_7} \frac{\lambda_2}{m_c^2} \simeq 2.5\%. \quad (5)$$

Here we used $C_2 = 1.11$, $C_7 = -0.32$, $\lambda_2 = 0.12 \text{ GeV}^2$, and $m_c = 1.4 \text{ GeV}$. This result is an order of magnitude larger than the perturbative estimate of the $O_2 O_7$ contribution (which contains a gluon in the final state). The $n = 1$ matrix element vanishes due to the equations of motion¹⁰. The $n > 1$ terms in Eq. (4) depend on an infinite series of unknown matrix elements. Estimating $\langle \bar{B}(v) | \bar{b} \Gamma^{\alpha\beta}(\hat{q}, v) (iD_{\mu_1} \dots iD_{\mu_n} g_s G_{\alpha\beta}) b | \bar{B}(v) \rangle / (2m_B) \sim (\Lambda_{\text{QCD}})^{n+2}$, the $n > 1$ terms are “suppressed” compared to the $n = 0$ term only by powers of $m_b \Lambda_{\text{QCD}} / m_c^2$. (A different estimate was given by Grant *et al.*¹⁰)

In the limit where m_c is fixed and $m_b \rightarrow \infty$, the higher order terms in Eq. (4) become successively more important and the expansion we have made is inappropriate. (The whole contribution in Eq. (3) is still suppressed by $\Lambda_{\text{QCD}} / m_b$.) In the limit where m_b / m_c is fixed and $m_{c,b} \rightarrow \infty$, the $n = 0$ result dominates the sum in Eq. (4) as the $n > 1$ terms are suppressed by powers of $\Lambda_{\text{QCD}} / m_c$. In the physical world, $m_b \Lambda_{\text{QCD}} / m_c^2 \sim 0.6$ is of order unity. As the coefficients of the operators in Eq. (4) are already small for small n , and decrease asymptotically as $3\sqrt{\pi} / (2^{n+1} n^{3/2})$, the $n > 1$ terms probably do not introduce an uncertainty larger than the size of the leading $n = 0$ term. Nonperturbative effects from the $O_1 O_7$ interference are expected to be smaller.

Consider next the contribution of $(C_1 O_1 + C_2 O_2)^2$. Diagrams like that in Fig. 1 should give a smaller result than for the interference of O_2 with O_7 (of order $\Lambda_{\text{QCD}}^4 / m_c^4$ instead of $\Lambda_{\text{QCD}}^2 / m_c^2$). But in this case there is a contribution to the $\bar{B} \rightarrow X_s \gamma$ decay rate from $\bar{B} \rightarrow X_s J/\psi$ followed by $J/\psi \rightarrow \gamma X$,

which is much larger than the perturbative calculation of the effect of $(C_1 O_1 + C_2 O_2)^2$. The combined branching ratio for this process is about as large as the total $\bar{B} \rightarrow X_s \gamma$ decay rate: of order 10^{-4} . This might not present a serious difficulty for the comparison of experiment with theory, since $\bar{B} \rightarrow X_s J/\psi$ followed by $J/\psi \rightarrow \gamma X$ does not favor hard photons. Moreover, it can be treated as a background and subtracted away. However, if such a subtraction is made, is it not double counting to include the perturbative result for the $(C_1 O_1 + C_2 O_2)^2$ contribution into the theoretical prediction? In any event, $B \rightarrow X_s J/\psi$ followed by $J/\psi \rightarrow \gamma X$ is a long distance contribution, while the charm quarks are far off-shell when nonperturbative effects suppressed by powers of Λ_{QCD}/m_c are calculable. Further work on these issues is warranted.

In summary, the nonperturbative contribution to the matrix element of $C_7 O_7$, of order $\Lambda_{\text{QCD}}^2/m_b^2$, is about -3% ³ with small uncertainty. Although there is no OPE for the contribution of photon coupling to light quarks, such effects give less than five percent uncertainty for hard photons. For the contribution of photon coupling to charm quarks, there are nonperturbative effects of order $\Lambda_{\text{QCD}}^2/m_c^2$, whose magnitude is 2.5%, with a similar uncertainty. However, it is possible that larger nonperturbative effects may exist. For comparison with the present data¹ which focuses on the region $E_\gamma \geq 2.2 \text{ GeV}$, the largest theoretical uncertainty is due to the contribution of higher dimension operators to T_{77} which become important in the end-point region. This uncertainty would be substantially smaller if the photon energy cut were reduced.

I thank Lisa Randall and Mark Wise for collaboration. This work was supported by the U.S. Dept. of Energy under grant no. DE-FG03-92-ER 40701.

References

1. CLEO Collaboration, M.S. Alam *et al.*, *Phys. Rev. Lett.* **74**, 2885 (1995).
2. T. Hurth, these proceedings, and references therein.
3. A.F. Falk *et al.*, *Phys. Rev. D* **49**, 3367 (1994);
I. Bigi *et al.*, in *The Fermilab Meeting*, Proc. of the Annual Meeting of the DPS of the APS, ed. C. Albright *et al.* (World Scientific, 1993).
4. E. Witten, *Nucl. Phys. B* **120**, 189 (1977); C.H. Llewellyn Smith, *Phys. Lett. B* **79**, 83 (1978); K. Koller *et al.*, *Z. Phys. C* **2**, 197 (1979).
5. N. Pott, *Phys. Rev. D* **54**, 938 (1996); C. Greub *et al.*, *ibid.* 3350.
6. A. Kapustin *et al.*, *Phys. Lett. B* **357**, 653 (1995).
7. J.F. Owens, *Rev. Mod. Phys.* **59**, 465 (1987); and references therein.
8. M.B. Voloshin, hep-ph/9612483.
9. Z. Ligeti *et al.*, hep-ph/9702322, to appear in *Phys. Lett. B*.
10. A.K. Grant *et al.*, hep-ph/9702380.